

Gene Section

Review

PIM1 (pim-1 oncogene)

Sai-Ching Jim Yeung

The University of Texas M. D. Anderson Cancer Center, Department of General Internal Medicine, Ambulatory Treatment and Emergency Care, Department of Endocrine Neoplasia and Hormonal Disorders, 1515 Holcombe Boulevard, Unit 437, Houston, Texas 77030, USA (SCJY)

Published in Atlas Database: April 2013

Online updated version : <http://AtlasGeneticsOncology.org/Genes/PIM1ID261.html>
DOI: 10.4267/2042/51537

This article is an update of :

Huret JL. PIM1 pim-1 oncogene. *Atlas Genet Cytogenet Oncol Haematol* 2000;4(4):183-184.

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 2013 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

Identity

Other names: PIM

HGNC (Hugo): PIM1

Location: 6p21.2

DNA/RNA

Description

PIM1 is a single gene with 5 introns and 6 exons that span 5 kb of DNA in the human genome (Meeker et al., 1987).

The gene starts at 37137922 and ends at 37143204 base pairs from pter. It is highly conserved evolutionarily across species.

Transcription

The mRNA sequence is 2,6 kb in length with a 941 bp coding region.

Protein

Description

Size: 313 amino acids; molecular weight: 36 kDa.

The PIM1 gene has six exons, but there are two isoforms of PIM1 protein, 34 kDa and 44 kDa, due to protein synthesis using alternative sites of translation initiation (Saris et al., 1991).

Both proteins show comparable kinase activities in vitro, but the 44 kDa isoform contains an N-terminal proline-rich motif that binds the ETK SH3 domain, and is recruited to the plasma membrane (Xie et al., 2006).

Expression

The kinase activity of all the PIM proteins is constitutively active, and there are no regulatory domains in the amino acid sequences of the PIM proteins.

Thus, unlike other kinases that are regulated by phosphorylation or binding to the plasma membrane, the activity of PIM1 is regulated primarily by transcription, translation and proteosomal degradation (Amaravadi and Thompson, 2005).

The gene expression of PIM1 is increased by various cytokines, mitogens and hormones such as G-CSF, GM-CSF, erythropoietin, interleukins, Con A, PMA, interferons, and prolactin (Wang et al., 2001; Hogan et al., 2008; White, 2003).

These factors act through the JAK/STAT pathway. The upregulation of PIM1 gene expression results from the binding site of STAT3 or STAT5 to the PIM1 gene promoter, and the ISFR/GAS-sequence (IFN- γ activation sequence) is an important binding site (Block et al., 2012; Matikainen et al., 1999; Yip-Schneider et al., 1995).

PIM1 phosphorylates and stabilizes SOCS proteins (suppressor of cytokine signaling) to provide negative feedback regulation of the JAK/STAT pathway (Peltola et al., 2004).

NF- κ B, as a downstream transcription factors, could also activate PIM1. In solid tumors, hypoxia would induce the PIM1 expression, independently of HIF1 α and by Krueppel-like factor 5 (KLF5) upon DNA damage (Chen et al., 2009a). ERG, as a transcription factors, also plays a role in PIM1 expression in the initial stages of prostate carcinogenesis (Magistroni et al., 2011).

Because of multiple copies of AUUU(A) motifs in the 3'UTR and GC-rich regions in the 5'UTR, mRNA of PIM genes are short lived (Wang et al., 2005).

Translation of PIM1 seems to be cap-dependent, and overexpression of eIF4E would increase PIM1 protein level (Hoover et al., 1997).

PIM RNA transcripts are regulatory targets of different miRNAs such as microRNAs miR1, miR-210, miR-33a, and miR328, implicating another layer of PIM expression regulation (Eiring et al., 2010; Huang et al., 2009; Nasser et al., 2008; Thomas et al., 2012).

At the post-translational level, the short half-life of PIM1 is primarily regulated by proteasomal degradation. PIM1 protein can be stabilized by the binding to HSP90 (Mizuno et al., 2001).

On the other hand, binding to HSP70 would induce the ubiquitylation of PIM1 and proteasomal degradation (Shay et al., 2005).

Also, in hypoxia, ubiquitin-mediated proteasomal degradation of PIM is prevented by HSP90 (Mizuno et al., 2001). PIM1 protein stability is further regulated by its phosphorylation status. PIM1 is able to autophosphorylate (Bullock et al., 2005). Phosphorylation by itself and/or other unknown kinases is important for PIM1 protein stability and function because PP2A phosphatase negatively regulates PIM1 stability (Losman et al., 2003).

Localisation

PIM1 is highly expressed in lymphoid and hematopoietic tissues (bone marrow, thymus, spleen, and fetal liver) (Eichmann et al., 2000) as well as in some non-hematopoietic tissues (e.g., hippocampus, oral epithelium, and the prostate gland). In some myeloid and lymphoid leukemia cell lines, prostate cancer cell lines, and also HeLa cells, PIM have also been detected. The PIM1 protein can be detected subcellularly in the cytoplasm and the nucleus.

Function

PIM1 phosphorylates a large subset of cellular substrates and thus regulates many different cellular processes such as cell cycle progression, cellular division, differentiation and apoptosis.

One of the cell-cycle-related targets of PIM1 is p21^{waf1} (Wang et al., 2002; Zhang et al., 2007). By phosphorylating the CDK inhibitor p21^{waf1} on T145, PIM1 lead to the nuclear export and inactivation of p21^{waf1}. Phosphorylation of another CDK inhibitor p27^{Kip1} at Thr157 and Thr198 would induce its proteasomal degradation and cell cycle progression.

Moreover, PIM1 seems to phosphorylate and inactivate the transcription factors of p27^{Kip1},

FoxO1a and FoxO3a (Morishita et al., 2008). Another mechanism of p27^{Kip1} regulation is the phosphorylation of SKP2 at Thr417, which control its stability and ability to degrade p27 (Cen et al., 2010). Additionally,

the phosphorylation of Cdc25A and Cdc25C would induce G1/S and G2/M transition, respectively (Bachmann et al., 2004).

PIM1 also implicate in mitosis promotion by interacts with dynein/dynactin and HP1 β (Magnuson et al., 2010).

Moreover, PIM1 is involved in genomic instability. By interaction with NuMA (nuclear mitotic apparatus protein), overexpression of PIM1 causes the loss of checkpoint control (Bhattacharya et al., 2002).

Consequently, cells with abnormal mitotic spindles are not arrested in mitosis, producing polyploid and multinucleated daughter cells.

PIM1 can also act as an oncogenic survival factor because of its function in blocking apoptotic cell death. It is consensus that phosphorylation of BAD at S112 would induce its proteasomal degradation and thus shifts the apoptosis threshold (Peltola et al., 2004).

The proapoptotic activity of ASK1 and PRAS40 would also be impaired by PIM1 phosphorylation (Gu et al., 2009; Zhang et al., 2009).

Through inactivation of ASK1 and subsequently less phosphorylation of the stress kinases JNK and p38, caspase-3 activation would be less and thus reduce cell death.

Moreover, the block of MDM2 and p53 degradation by PIM1 may induce senescence in embryonic fibroblasts and cancer cells (Hogan et al., 2008).

When bound to MYC at the E-box, PIM1 would participate in the complex's phosphorylation of histone H3 at S10 and thus participate in the stimulation of transcription of a specific subset of MYC-dependent genes (Zippo et al., 2007).

Additionally, PIM1 influences the activity of a number of transcriptional regulators, such as HP-1, PAP-1, TFAF2/SNX6, NFATc1, p100, RUNX, SOCS1, RelA/p65 and c-Myb (Bhattacharya et al., 2002; Evans and Fox, 2007; Ishibashi et al., 2001; Kim et al., 2010; Rainio et al., 2002; Winn et al., 2003).

PIM kinases also phosphorylate 4E-BP1, inhibiting its binding to eIF-4E.

Since eIF-4E is a rate-limiting factor in protein synthesis, PIM kinases may also regulate 5' cap-dependent translation (Beharry et al., 2011).

Implicated in

t(3;6)(q27;p21.2) in diffuse large B-cell lymphoma (DLCL); chimeric BCL6 / PIM1

Note

Only 1 case to date.

Hybrid/Mutated gene

5' PIM1 fused to 3' BCL6; the substitution of the promoter of BCL6 causes deregulation of BCL6.

Myeloid and lymphoid leukemias and other lymphomas

Oncogenesis

PIM1 induces anti-apoptotic oncogenes such as BCR/ABL, FLT2, CBL or JAK2 (Adam et al., 2006; Mizuki et al., 2003; Naramura et al., 2011; Nieborowska-Skorska et al., 2002; Wernig et al., 2008). PIM1 mRNA is upregulated in acute myeloid leukemia (AML) along with MLL gene rearrangements, e.g., MLL/AF9 or MLL/ENL (Chen et al., 2008). The reason for PIM1 levels increase seems to be the constitutive activation of FLT3 or Hoxa9 (Hu et al., 2007). Additionally, PIM1 also involve in the several B-cell developmental disorders that are related to Kaposi sarcoma associated herpesvirus (KSHV) or the Epstein-Barr virus (EBV) (Bajaj et al., 2006; Cheng et al., 2009).

Prostate cancer

Prognosis

In more half of the prostate cancer samples, PIM1 is showed a relatively overexpression compared to benign lesions and the expression elevation correlated with a poor therapeutic outcome (Dhanasekaran et al., 2001).

Oncogenesis

In mouse model, the synergistic effects of PIM1 and MYC showed obvious co-regulation in prostate cancer. The molecular mechanism for the oncogenic activity might because PIM1 phosphorylation of c-MYC would increase its half-life and also because PIM1 enhancement of transcriptional activity of c-MYC (Chen et al., 2009b; Mumenthaler et al., 2009). Moreover, PIM1 kinase is related to chemoresistance in prostate cancer cells, which is related to relatively aggressive or hormone-refractory prostate cancers. The high level of expression of PIM1 in high grade prostate intraepithelial neoplasia may indicate a role of PIM1 in the early prostate carcinogenesis. PIM1 is also found to be upregulated in patients undergoing androgen ablation therapy (van der Poel et al., 2010).

Pancreatic cancer

Note

Hypoxia-promoted genetic instability

Oncogenesis

PIM1 increases in hypoxic condition, independently of HIF-1 α (Reiser-Erkan et al., 2008). It is now proposed as a prognostic marker. Increase in PIM1 expression may partly account for resistance to chemotherapy.

Sporadic malignant tumors

Oncogenesis

Overexpression of PIM1 is founded in gastric carcinoma, squamous cell carcinoma, colorectal carcinoma, liver carcinoma (Shah et al., 2008), liposarcoma (Nga et al., 2010), and bladder cancer (Guo et al., 2010).

References

- Selten G, Cuypers HT, Boelens W, Robanus-Maandag E, Verbeek J, Domen J, van Beveren C, Berns A. The primary structure of the putative oncogene *pim-1* shows extensive homology with protein kinases. *Cell*. 1986 Aug 15;46(4):603-11
- Meeker TC, Nagarajan L, ar-Rushdi A, Croce CM. Cloning and characterization of the human PIM-1 gene: a putative oncogene related to the protein kinases. *J Cell Biochem*. 1987 Oct;35(2):105-12
- Saris CJ, Domen J, Berns A. The *pim-1* oncogene encodes two related protein-serine/threonine kinases by alternative initiation at AUG and CUG. *EMBO J*. 1991 Mar;10(3):655-64
- Yip-Schneider MT, Horie M, Broxmeyer HE. Transcriptional induction of *pim-1* protein kinase gene expression by interferon gamma and posttranscriptional effects on costimulation with steel factor. *Blood*. 1995 Jun 15;85(12):3494-502
- Hoover DS, Wingett DG, Zhang J, Reeves R, Magnuson NS. Pim-1 protein expression is regulated by its 5'-untranslated region and translation initiation factor eIF-4E. *Cell Growth Differ*. 1997 Dec;8(12):1371-80
- Mochizuki T, Kitanaka C, Noguchi K, Sugiyama A, Kagaya S, Chi S, Asai A, Kuchino Y. Pim-1 kinase stimulates c-Myc-mediated death signaling upstream of caspase-3 (CPP32)-like protease activation. *Oncogene*. 1997 Sep 18;15(12):1471-80
- Levenson JD, Koskinen PJ, Orrico FC, Rainio EM, Jalkanen KJ, Dash AB, Eisenman RN, Ness SA. Pim-1 kinase and p100 cooperate to enhance c-Myb activity. *Mol Cell*. 1998 Oct;2(4):417-25
- Matikainen S, Sareneva T, Ronni T, Lehtonen A, Koskinen PJ, Julkunen I. Interferon-alpha activates multiple STAT proteins and upregulates proliferation-associated IL-2R α , c-myc, and *pim-1* genes in human T cells. *Blood*. 1999 Mar 15;93(6):1980-91
- Mochizuki T, Kitanaka C, Noguchi K, Muramatsu T, Asai A, Kuchino Y. Physical and functional interactions between Pim-1 kinase and Cdc25A phosphatase. Implications for the Pim-1-mediated activation of the c-Myc signaling pathway. *J Biol Chem*. 1999 Jun 25;274(26):18659-66
- Shirogane T, Fukada T, Muller JM, Shima DT, Hibi M, Hirano T. Synergistic roles for Pim-1 and c-Myc in STAT3-mediated cell cycle progression and antiapoptosis. *Immunity*. 1999 Dec;11(6):709-19
- Yoshida S, Kaneita Y, Aoki Y, Seto M, Mori S, Moriyama M. Identification of heterologous translocation partner genes fused to the BCL6 gene in diffuse large B-cell lymphomas: 5'-RACE and LA - PCR analyses of biopsy samples. *Oncogene*. 1999 Dec 23;18(56):7994-9
- Eichmann A, Yuan L, Bréant C, Alitalo K, Koskinen PJ. Developmental expression of *pim* kinases suggests functions also outside of the hematopoietic system. *Oncogene*. 2000 Feb 24;19(9):1215-24
- Koike N, Maita H, Taira T, Ariga H, Iguchi-Ariga SM. Identification of heterochromatin protein 1 (HP1) as a phosphorylation target by Pim-1 kinase and the effect of phosphorylation on the transcriptional repression function of HP1(1). *FEBS Lett*. 2000 Feb 4;467(1):17-21
- Dhanasekaran SM, Barrette TR, Ghosh D, Shah R, Varambally S, Kurachi K, Pienta KJ, Rubin MA, Chinnaiyan AM. Delineation of prognostic biomarkers in prostate cancer. *Nature*. 2001 Aug 23;412(6849):822-6

- Ishibashi Y, Maita H, Yano M, Koike N, Tamai K, Ariga H, Iguchi-Ariga SM. Pim-1 translocates sorting nexin 6/TRAFF4-associated factor 2 from cytoplasm to nucleus. *FEBS Lett*. 2001 Sep 28;506(1):33-8
- Mizuno K, Shirogane T, Shinohara A, Iwamatsu A, Hibi M, Hirano T. Regulation of Pim-1 by Hsp90. *Biochem Biophys Res Commun*. 2001 Mar 2;281(3):663-9
- Wang Z, Bhattacharya N, Weaver M, Petersen K, Meyer M, Gapter L, Magnuson NS. Pim-1: a serine/threonine kinase with a role in cell survival, proliferation, differentiation and tumorigenesis. *J Vet Sci*. 2001 Dec;2(3):167-79
- Bhattacharya N, Wang Z, Davitt C, McKenzie IF, Xing PX, Magnuson NS. Pim-1 associates with protein complexes necessary for mitosis. *Chromosoma*. 2002 Jul;111(2):80-95
- Nieborowska-Skorska M, Hoser G, Kossev P, Wasik MA, Skorski T. Complementary functions of the antiapoptotic protein A1 and serine/threonine kinase pim-1 in the BCR/ABL-mediated leukemogenesis. *Blood*. 2002 Jun 15;99(12):4531-9
- Rainio EM, Sandholm J, Koskinen PJ. Cutting edge: Transcriptional activity of NFATc1 is enhanced by the Pim-1 kinase. *J Immunol*. 2002 Feb 15;168(4):1524-7
- Wang Z, Bhattacharya N, Mixter PF, Wei W, Sedivy J, Magnuson NS. Phosphorylation of the cell cycle inhibitor p21Cip1/WAF1 by Pim-1 kinase. *Biochim Biophys Acta*. 2002 Dec 16;1593(1):45-55
- Losman JA, Chen XP, Vuong BQ, Fay S, Rothman PB. Protein phosphatase 2A regulates the stability of Pim protein kinases. *J Biol Chem*. 2003 Feb 14;278(7):4800-5
- Mizuki M, Schwable J, Steur C, Choudhary C, Agrawal S, Sargin B, Steffen B, Matsumura I, Kanakura Y, Böhmer FD, Müller-Tidow C, Berdel WE, Serve H. Suppression of myeloid transcription factors and induction of STAT response genes by AML-specific Flt3 mutations. *Blood*. 2003 Apr 15;101(8):3164-73
- White E. The pims and outs of survival signaling: role for the Pim-2 protein kinase in the suppression of apoptosis by cytokines. *Genes Dev*. 2003 Aug 1;17(15):1813-6
- Winn LM, Lei W, Ness SA. Pim-1 phosphorylates the DNA binding domain of c-Myb. *Cell Cycle*. 2003 May-Jun;2(3):258-62
- Bachmann M, Hennemann H, Xing PX, Hoffmann I, Möröy T. The oncogenic serine/threonine kinase Pim-1 phosphorylates and inhibits the activity of Cdc25C-associated kinase 1 (C-TAK1): a novel role for Pim-1 at the G2/M cell cycle checkpoint. *J Biol Chem*. 2004 Nov 12;279(46):48319-28
- Kim O, Jiang T, Xie Y, Guo Z, Chen H, Qiu Y. Synergism of cytoplasmic kinases in IL6-induced ligand-independent activation of androgen receptor in prostate cancer cells. *Oncogene*. 2004 Mar 11;23(10):1838-44
- Peltola KJ, Pauku K, Aho TL, Ruuska M, Silvennoinen O, Koskinen PJ. Pim-1 kinase inhibits STAT5-dependent transcription via its interactions with SOCS1 and SOCS3. *Blood*. 2004 May 15;103(10):3744-50
- Amaravadi R, Thompson CB. The survival kinases Akt and Pim as potential pharmacological targets. *J Clin Invest*. 2005 Oct;115(10):2618-24
- Bullock AN, Debreczeni JE, Fedorov OY, Nelson A, Marsden BD, Knapp S. Structural basis of inhibitor specificity of the human protooncogene proviral insertion site in moloney murine leukemia virus (PIM-1) kinase. *J Med Chem*. 2005 Dec 1;48(24):7604-14
- Shay KP, Wang Z, Xing PX, McKenzie IF, Magnuson NS. Pim-1 kinase stability is regulated by heat shock proteins and the ubiquitin-proteasome pathway. *Mol Cancer Res*. 2005 Mar;3(3):170-81
- Wang Z, Weaver M, Magnuson NS. Cryptic promoter activity in the DNA sequence corresponding to the pim-1 5'-UTR. *Nucleic Acids Res*. 2005;33(7):2248-58
- Adam M, Pogacic V, Bendit M, Chappuis R, Nawijn MC, Duyster J, Fox CJ, Thompson CB, Cools J, Schwaller J. Targeting PIM kinases impairs survival of hematopoietic cells transformed by kinase inhibitor-sensitive and kinase inhibitor-resistant forms of Fms-like tyrosine kinase 3 and BCR/ABL. *Cancer Res*. 2006 Apr 1;66(7):3828-35
- Bajaj BG, Verma SC, Lan K, Cotter MA, Woodman ZL, Robertson ES. KSHV encoded LANA upregulates Pim-1 and is a substrate for its kinase activity. *Virology*. 2006 Jul 20;351(1):18-28
- Xie Y, Xu K, Dai B, Guo Z, Jiang T, Chen H, Qiu Y. The 44 kDa Pim-1 kinase directly interacts with tyrosine kinase Etk/BMX and protects human prostate cancer cells from apoptosis induced by chemotherapeutic drugs. *Oncogene*. 2006 Jan 5;25(1):70-8
- Evans KE, Fox SW. Interleukin-10 inhibits osteoclastogenesis by reducing NFATc1 expression and preventing its translocation to the nucleus. *BMC Cell Biol*. 2007 Jan 19;8:4
- Hu YL, Passequé E, Fong S, Largman C, Lawrence HJ. Evidence that the Pim1 kinase gene is a direct target of HOXA9. *Blood*. 2007 Jun 1;109(11):4732-8
- Zhang Y, Wang Z, Magnuson NS. Pim-1 kinase-dependent phosphorylation of p21Cip1/WAF1 regulates its stability and cellular localization in H1299 cells. *Mol Cancer Res*. 2007 Sep;5(9):909-22
- Zippo A, De Robertis A, Serafini R, Oliviero S. PIM1-dependent phosphorylation of histone H3 at serine 10 is required for MYC-dependent transcriptional activation and oncogenic transformation. *Nat Cell Biol*. 2007 Aug;9(8):932-44
- Chen W, Kumar AR, Hudson WA, Li Q, Wu B, Staggs RA, Lund EA, Sam TN, Kersey JH. Malignant transformation initiated by Mll-AF9: gene dosage and critical target cells. *Cancer Cell*. 2008 May;13(5):432-40
- Hogan C, Hutchison C, Marcar L, Milne D, Saville M, Goodlad J, Kernohan N, Meek D. Elevated levels of oncogenic protein kinase Pim-1 induce the p53 pathway in cultured cells and correlate with increased Mdm2 in mantle cell lymphoma. *J Biol Chem*. 2008 Jun 27;283(26):18012-23
- Morishita D, Katayama R, Sekimizu K, Tsuruo T, Fujita N. Pim kinases promote cell cycle progression by phosphorylating and down-regulating p27Kip1 at the transcriptional and posttranscriptional levels. *Cancer Res*. 2008 Jul 1;68(13):5076-85
- Nasser MW, Datta J, Nuovo G, Kutay H, Motiwala T, Majumder S, Wang B, Suster S, Jacob ST, Ghoshal K. Down-regulation of micro-RNA-1 (miR-1) in lung cancer. Suppression of tumorigenic property of lung cancer cells and their sensitization to doxorubicin-induced apoptosis by miR-1. *J Biol Chem*. 2008 Nov 28;283(48):33394-405
- Reiser-Erkan C, Erkan M, Pan Z, Bekasi S, Giese NA, Streit S, Michalski CW, Friess H, Kleeff J. Hypoxia-inducible proto-oncogene Pim-1 is a prognostic marker in pancreatic ductal adenocarcinoma. *Cancer Biol Ther*. 2008 Sep;7(9):1352-9
- Shah N, Pang B, Yeoh KG, Thorn S, Chen CS, Lilly MB, Salto-Tellez M. Potential roles for the PIM1 kinase in human cancer - a molecular and therapeutic appraisal. *Eur J Cancer*. 2008 Oct;44(15):2144-51
- Wernig G, Gonneville JR, Crowley BJ, Rodrigues MS, Reddy MM, Hudon HE, Walz C, Reiter A, Podar K, Royer Y, Constantinescu SN, Tomasson MH, Griffin JD, Gilliland DG,

- Sattler M. The Jak2V617F oncogene associated with myeloproliferative diseases requires a functional FERM domain for transformation and for expression of the Myc and Pim proto-oncogenes. *Blood*. 2008 Apr 1;111(7):3751-9
- Chen J, Kobayashi M, Darmanin S, Qiao Y, Gully C, Zhao R, Kondo S, Wang H, Wang H, Yeung SC, Lee MH. Hypoxia-mediated up-regulation of Pim-1 contributes to solid tumor formation. *Am J Pathol*. 2009a Jul;175(1):400-11
- Chen J, Kobayashi M, Darmanin S, Qiao Y, Gully C, Zhao R, Yeung SC, Lee MH. Pim-1 plays a pivotal role in hypoxia-induced chemoresistance. *Oncogene*. 2009b Jul 16;28(28):2581-92
- Cheng F, Weidner-Glunde M, Varjosalo M, Rainio EM, Lehtonen A, Schulz TF, Koskinen PJ, Taipale J, Ojala PM. KSHV reactivation from latency requires Pim-1 and Pim-3 kinases to inactivate the latency-associated nuclear antigen LANA. *PLoS Pathog*. 2009 Mar;5(3):e1000324
- Gu JJ, Wang Z, Reeves R, Magnuson NS. PIM1 phosphorylates and negatively regulates ASK1-mediated apoptosis. *Oncogene*. 2009 Dec 3;28(48):4261-71
- Huang X, Ding L, Bennewith KL, Tong RT, Welford SM, Ang KK, Story M, Le QT, Giaccia AJ. Hypoxia-inducible mir-210 regulates normoxic gene expression involved in tumor initiation. *Mol Cell*. 2009 Sep 24;35(6):856-67
- Mumenthaler SM, Ng PY, Hodge A, Bearss D, Berk G, Kanekal S, Redkar S, Taverna P, Agus DB, Jain A. Pharmacologic inhibition of Pim kinases alters prostate cancer cell growth and resensitizes chemoresistant cells to taxanes. *Mol Cancer Ther*. 2009 Oct;8(10):2882-93
- Zhang F, Beharry ZM, Harris TE, Lilly MB, Smith CD, Mahajan S, Kraft AS. PIM1 protein kinase regulates PRAS40 phosphorylation and mTOR activity in FDCP1 cells. *Cancer Biol Ther*. 2009 May;8(9):846-53
- Cen B, Mahajan S, Zemskova M, Beharry Z, Lin YW, Cramer SD, Lilly MB, Kraft AS. Regulation of Skp2 levels by the Pim-1 protein kinase. *J Biol Chem*. 2010 Sep 17;285(38):29128-37
- Eiring AM, Harb JG, Neviani P, Garton C, Oaks JJ, Spizzo R, Liu S, Schwind S, Santhanam R, Hickey CJ, Becker H, Chandler JC, Andino R, Cortes J, Hokland P, Huettner CS, Bhatia R, Roy DC, Liebhaber SA, Caligiuri MA, Marcucci G, Garzon R, Croce CM, Calin GA, Perrotti D. miR-328 functions as an RNA decoy to modulate hnRNP E2 regulation of mRNA translation in leukemic blasts. *Cell*. 2010 Mar 5;140(5):652-65
- Guo S, Mao X, Chen J, Huang B, Jin C, Xu Z, Qiu S. Overexpression of Pim-1 in bladder cancer. *J Exp Clin Cancer Res*. 2010 Dec 11;29:161
- Kim K, Kim JH, Youn BU, Jin HM, Kim N. Pim-1 regulates RANKL-induced osteoclastogenesis via NF- κ B activation and NFATc1 induction. *J Immunol*. 2010 Dec 15;185(12):7460-6
- Magnuson NS, Wang Z, Ding G, Reeves R. Why target PIM1 for cancer diagnosis and treatment? *Future Oncol*. 2010 Sep;6(9):1461-78
- Nga ME, Swe NN, Chen KT, Shen L, Lilly MB, Chan SP, Salto-Tellez M, Das K. PIM-1 kinase expression in adipocytic neoplasms: diagnostic and biological implications. *Int J Exp Pathol*. 2010 Feb;91(1):34-43
- van der Poel HG, Zevenhoven J, Bergman AM. Pim1 regulates androgen-dependent survival signaling in prostate cancer cells. *Urol Int*. 2010;84(2):212-20
- Beharry Z, Mahajan S, Zemskova M, Lin YW, Tholanikunnel BG, Xia Z, Smith CD, Kraft AS. The Pim protein kinases regulate energy metabolism and cell growth. *Proc Natl Acad Sci U S A*. 2011 Jan 11;108(2):528-33
- Magistrini V, Mologni L, Sanselicio S, Reid JF, Redaelli S, Piazza R, Viltadi M, Bovo G, Strada G, Grasso M, Gariboldi M, Gambacorti-Passerini C. ERG deregulation induces PIM1 over-expression and aneuploidy in prostate epithelial cells. *PLoS One*. 2011;6(11):e28162
- Naramura M, Nadeau S, Mohapatra B, Ahmad G, Mukhopadhyay C, Sattler M, Raja SM, Natarajan A, Band V, Band H. Mutant Cbl proteins as oncogenic drivers in myeloproliferative disorders. *Oncotarget*. 2011 Mar;2(3):245-50
- Block KM, Hanke NT, Maine EA, Baker AF. IL-6 stimulates STAT3 and Pim-1 kinase in pancreatic cancer cell lines. *Pancreas*. 2012 Jul;41(5):773-81
- Thomas M, Lange-Grünweller K, Weirauch U, Gutsch D, Aigner A, Grünweller A, Hartmann RK. The proto-oncogene Pim-1 is a target of miR-33a. *Oncogene*. 2012 Feb 16;31(7):918-28

This article should be referenced as such:

Yeung SCJ. PIM1 (*pim-1* oncogene). *Atlas Genet Cytogenet Oncol Haematol*. 2013; 17(10):704-708.
